

REFERENCES

- Akyol, O., Kinacigil, H.T. & Sevik, R. (2007). Longline fishery and length-weight relationships for selected fish species in Gokova Bay (Aegean Sea, Turkey). *International Journal of Natural and Engineering Sciences*, 1, 1-4.
- Biçer, D., Yıldız, T., Uzer, U., Karakulak, F.S. (2020). Last dalians (stationary uncovered pound nets) deployed in the Istanbul Strait: Catch composition, catch per unit effort, and ecological indexes. *Ege Journal of Fisheries and Aquatic Sciences* 37(2), 125-133.
- Bok, T.D., Gokturk, D., Kahraman, A.E., Alicli, T.Z., Acun, T. & Ateş, C. (2011). Length-weight relationships of 34 fish species from the Sea of Marmara, Turkey. *Journal of Animal and Veterinary Advances*, 10 (23): 3037-3042. DOI: [10.3923/javaa.2011.3037.3042](https://doi.org/10.3923/javaa.2011.3037.3042)
- Cengiz, Ö. (2013). Length-weight relationships of 22 fish species from the Gallipoli Peninsula and Dardanelles (northeastern Mediterranean, Turkey). *Turkish Journal of Zoology*, 37: 419-422. DOI: [10.3906/zoo-1209-30](https://doi.org/10.3906/zoo-1209-30)
- Ceyhan, T., Akyol, O. & Erdem, M. (2009). Length-weight relationships of fishes from Gokova Bay, Turkey (Aegean Sea). *Turkish Journal of Zoology*, 33: 69-72.
- Çolakoğlu, S., Tokaç, A., İşmen, A. & Yurdusev, H. (2015). Catch composition of set net (fixed stake trap) fisheries in the coastal waters of Saros Bay, North Aegean Sea. *Ege Journal of Fisheries Aquatic Science* 32(2): 53-58. DOI: [10.12714/egejfas.2015.32.2.02](https://doi.org/10.12714/egejfas.2015.32.2.02)
- Demirhan, S.A. & Can, M.F. (2007). Length-weight relationships for seven fish species from the southeastern Black Sea. *Journal of Applied Ichthyology*, 23: 282-283. DOI: [10.1111/j.14390426.2007.00835.x](https://doi.org/10.1111/j.14390426.2007.00835.x)
- Deveciyan, K. (2011). Türkiye'de Balık ve Balıkçılık. Aras Yayıncılık, İstanbul, 2011.
- Erguden, D., Turan, C. & Gurlek, M. (2009). Weight-length relationships for 20 lessepsian fish species caught by bottom trawl on the coast of Iskenderun Bay (NE Mediterranean Sea, Turkey). *Journal of Applied Ichthyology*, 25: 133-135. DOI: [10.1111/j.1439-0426.2008.01198](https://doi.org/10.1111/j.1439-0426.2008.01198)
- Filiz, H. & Bilge, B. (2004). Length-weight relationships of 24 fish species from the North Aegean Sea, Turkey. *Journal of Applied Ichthyology*, 20: 431-432. DOI: [10.1111/j.1439-0426.2004.00582.x](https://doi.org/10.1111/j.1439-0426.2004.00582.x)
- Froese, R. (2006). Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology*, 22: 241-253. DOI: [10.1111/j.1439-0426.2006.00805.x](https://doi.org/10.1111/j.1439-0426.2006.00805.x)
- Froese, R. & Pauly, D. (2013). FishBase. World Wide Web electronic publication. www.fishbase.org (10/2013).
- Gonçalves, J.M.S., Bentes, L., Lino, G., Almeida, J., Canário, A.V.M. & Erzini, K. (1997). Weight-length relationships for selected fish species of the small-scale demersal fisheries of the south and southwest coast of Portugal. *Fisheries Research*, 30, 251-256.
- Ilkyaz, T., Metin, G., Biçer, D. & Kinacigil, H.T. (2008). Length-weight relationship of 12 fish species from the Central Aegean Sea, Turkey. *Journal of Applied Ichthyology*, 24, 699-702. DOI: [10.1111/j.1439-0426.2008.01167.x](https://doi.org/10.1111/j.1439-0426.2008.01167.x)
- İsmeň, A., Ozen, O., Altınagac, U., Ozekinci, U. & Ayaz, A. (2007). Weight-length relationships of 63 fish species in Saros Bay, Turkey. *Journal of Applied Ichthyology*, 23, 707-708. DOI: [10.1111/j.1439-0426.2007.00872.x](https://doi.org/10.1111/j.1439-0426.2007.00872.x)
- Jobling, M. (2002). Environmental factors and rates of development and growth. In: Handbook of fish biology and fisheries (P.J.B. Hart & J.D. Reynolds eds). Pp. 97-122. Blackwell Science Ltd: Oxford, USA.
- Karakulak, F.S., Erk, H. & Bilgin, B. (2006). Length-weight relationships for 47 coastal fish species from the northern Aegean Sea, Turkey. *Journal of Applied Ichthyology*, 22, 274-278. DOI: [10.1111/j.1439-0426.2006.00736.x](https://doi.org/10.1111/j.1439-0426.2006.00736.x)
- Kasapoglu, N. & Duzgunes, E. (2014). Length-weight relationships of marine species caught by five gears from the Black Sea. *Mediterranean Marine Science*, 15, 95-100. DOI: [10.12681/mms.463](https://doi.org/10.12681/mms.463)
- Keskin, C. & Gaygusuz, O. (2010). Length-weight relationships of fishes in shallow waters of Erdek Bay (Sea of Marmara, Turkey). *IUFES Journal of Biology*, 69 (1), 25-32.
- Mater, S., Kaya, M. & Bilecenoğlu, M. (2007). Türkiye Deniz Balıkları Atlası (III. Baskı), Ege Üniversitesi Su Ürünleri Fakültesi Yayınları, No. 68, İzmir (In Turkish), 169p.
- Moutopoulos, D.K. & Stergiou, K.I. (2002). Length-weight and length-length relationships of fish species from the Aegean Sea (Greece). *Journal of Applied Ichthyology*, 18, 200-206. DOI: [10.1046/j.1439-0426.2002.00031.x](https://doi.org/10.1046/j.1439-0426.2002.00031.x)
- Ozaydin, O. & Taskavak, E. (2006). Length-weight relationships for 47 fish species from Izmir Bay, eastern Aegean Sea, Turkey). *Acta Adriatica*, 47, 211-216.
- Ozaydin, O., Uckun, M., Akalin, S., Leblebici, S. & Tosunoglu, Z. (2007). Length-weight relationships of fishes captured from Izmir Bay, Central Aegean Sea. *Journal of Applied Ichthyology*, 23, 695-696. DOI: [10.1111/j.1439-0426.2007.00853.x](https://doi.org/10.1111/j.1439-0426.2007.00853.x)
- Ricker, W.E. (1973). Linear regressions in fishery research. *Journal of the Fisheries Research Board of Canada*, 30, 409-434.
- Sangu, L., Akamca, E. & Akar, M. (2007). Weight-length relationships for 39 fish species from the north-eastern Mediterranean coast of Turkey. *Turkish Journal of Fisheries and Aquatic Sciences*, 7, 37-40.
- Silva, J.F., Ellis, J.R. & Ayers, R.A. (2013). Length-weight relationships of marine fish collected from around the British Isles. Science Series Technical Report n°150. Cefas: Lowestoft. 109 pp.
- StatSoft, Inc. (2007). STATISTICA (data analysis software system), version 8.0.
- Taskavak, E. & Bilecenoglu, M. (2001). Length-weight relationships for 18 lessepsian (Red Sea) immigrant fish species from the eastern Mediterranean coast of Turkey. *Journal of the Marine Biological Association of the United Kingdom*, 81, 895-896. DOI: [10.1017/S0025315401004805](https://doi.org/10.1017/S0025315401004805)
- Wootton, R.J. (1990). Ecology of Teleost Fish. Chapman and Hall, London. 404p.
- Yankova, M., Pavlov, D., Raykov, V., Michneva, V. & Radu, G. (2011). Length-Weight Relationships of Ten Fish Species from the Bulgarian Black Sea waters. *Turkish Journal of Zoology*, 35, 265-270.
- Yıldız, T., Zengin, M., Uzer, U., Akpınar, İ.Ö. & Karakulak, F.S. (2018). Length-weight relationships for 24 fish species collected from the western Black Sea (Turkey). *Cahiers de Biologie Marine*, 59,159-165. DOI: [10.21411/CBM.A.39436D05](https://doi.org/10.21411/CBM.A.39436D05)
- Zar, J.H. (1999). Biostatistical Analysis. 4th edition. Prentice Hall: Englewood Cliffs, New Jersey. 929 pp.