

On the first record of *Seirospora giraudyi* (Kützing) De Toni (Rhodophyceae, Ceramiales) from Turkey

Ergün Taşkın

Department of Biology, Faculty of Arts and Sciences, Celal Bayar University, Muradiye, Manisa, 45040, Turkey
E mail: ergun.taskin@bayar.edu.tr

In the paper entitled "Marine Flora of Sinop (Black Sea, Turkey)" by Aysel *et al.* (2004), in the E.U. Journal of Fisheries & Aquatic Sciences [21 (1-2): 59-68], *Seirospora giraudyi* (Kützing) De Toni (Rhodophyceae, Ceramiales) was reported as a new record for Turkey from the Sinop (Black Sea). But this species was reported before from the Iskenderun Gulf (Hatay, northeastern Mediterranean Sea) by Taşkın *et al.* (2000).

References

- Aysel, V., H. Erduğan, B. Dural-Tarakçı, E.Ş. Okudan, A. Şenkardeşler and F. Aysel. 2004. Marine Flora of Sinop (Black Sea, Turkey). E.U. Journal of Fisheries & Aquatic Sciences, 21 (1-2): 59 - 68.
- Taşkın, E., M. Öztürk, V. Aysel and O. Kurt. 2000. Specimens of Rhodophyta of Iskenderun Gulf (Hatay Shores / Turkey) and Ecologies. 19 May Univ. Faculty of Fisheries Engineering. "SINOP-2000" Symposium of Fisheries (20-22th Sept. 2000): 300-315.